Public Health District 1 Health Indicators

Panhandle Health District (Kootenai, Bonner, Benewah, Boundary and Shoshone Counties)

2006 Data

| Item | ldaho | District 1 |
|---|-------|------------|
| Maternal/Child | | |
| 1) % unintended pregnancies | 34.4 | 39.3 |
| 2a) % live births with adequate prenatal care in 2006 ¹ | 42.0 | 46.6 |
| 2b) % live births with intensive prenatal care in 2006 ¹ | 31.9 | 34.9 |
| 2c) % live births with adequate or intensive prenatal care in 2006 ¹ | 73.9 | 81.5 |
| 3a) % live births with adequate prenatal care in 2005 ¹ | 41.9 | 44.4 |
| 3b) % live births with intensive prenatal care in 2005 ¹ | 32.3 | 34.9 |
| 3c) % live births with adequate or intensive prenatal care in 2005 ¹ | 74.2 | 79.3 |
| 4) % live births with low birth weight ² | 6.9 | 5.6 |
| 5) % live births with tobacco use during pregnancy ³ | 12.7 | 21.6 |
| 6) % WIC participation ⁴ | 35.9 | 38.8 |
| 7) % prevalence of breastfeeding ⁵ | 88.4 | 89.0 |
| Adolescents | | |
| 1) Teen pregnancy rate (15-19) ⁶ | 46.5 | 42.4 |
| 2) Motor vehicle crash death rate, 2004-2006 (ages 15-19) ^{7, 9} | 27.0 | 25.3 |
| 3. Suicide rate, 2004-2006 (ages 10-18) 8,9 | 5.8 | 5.2 |
| Adults | | |
| 1) % without health care coverage | 18.8 | 20.6 |
| 2) % with no leisure time activity | 20.8 | 16.8 |
| 3) % overweight BMI ≥ 25) | 59.7 | 62.7 |
| 4) % diagnosed with diabetes | 6.8 | 7.0 |
| 5) % who smoke | 16.8 | 19.2 |
| 6) % who binge drink ¹⁰ | 14.8 | 17.4 |
| 7) % of females without breast cancer screening | 37.8 | 38.2 |
| 8) % of males without prostate cancer screening | 48.7 | 49.6 |
| 9) % who did not wear seatbelts | 26.8 | 22.2 |
| 10) Suicide rate, 2003-2005 (ages 65+) ^{8, 9} | 22.6 | 28.8 |
| 11) % with no dental visit in the past 12 months | 34.6 | 37.8 |

| Source | s: | | | |
|--------|----|--|--|--|
| | | | | |

Records and Health Statistics.

For Maternal/Child indicator 1: Pregnancy Risk Assessment System (PRATS), Bureau of Vital Records and Health Statistics. PRATS is a representative sample of resident women aged 18+ who gave birth in Idaho.

For Maternal/Child indicators 2-7, Adolescents indicators, and Adult indicator 10: Vital Statistics, Bureau of Vital Records and Health Statistics.

For Adult indicators 1-9, 11: Idaho Behavioral Risk Factor Surveillance System (BRFSS), Bureau of Vital Records and Health Statistics.

Notes:

- 1.In 2004, the Idaho birth certificate was revised. Beginning in 2004, Idaho prenatal care data are based on date of first prenatal care visit as reported in the mother's medical record. 2004 and 2005 APNCU data are not comparable with APNCU data based on month prenatal care began. The APNCU index classifies care as intensive, adequate, intermediate, inadequate, or no care by comparing the number of actual prenatal care visits that a woman was expected to receive, given the onset of prenatal care and the length of gestation. Percents based on records with stated data for APNCU calculation (date of first prenatal care visit, number of visits, and length of gestation). Data for 2004 have been corrected from the 2004 Health Indicators; 2004 data are comparable to 2005 data.
- Percents based on records with stated birth weight.
- Prior to 2004, tobacco use during pregnancy (yes or no) was reported on the Idaho birth certificate.
 Beginning in 2004, cigarette smoking before and during pregnancy is reported on the revised birth certificate.
 Data in 2004 and 2005 are not comparable to data prior to 2004.
- 4. The revised 2004 Idaho birth certificate includes a new variable for reporting if mother received WIC food for herself during this pregnancy (yes or no)?
- 5. The revised 2004 birth certificate includes a new variable for reporting if infant was being breastfed at birth or at the time the birth certificate was completed (yes or no)?
- 6. Pregnancies include live births, induced abortions, and reportable stillbirths (only those fetal deaths with a gestational period of 20+ weeks or which weight 350+ grams are required to be reportable under Idaho law.)
 Rate: Number of pregnancies per 1,000 females in corresponding age group. Rates are based on July 1, 2006 female population estimates from the National Center for Health Statistics, United States resident population from the Vintage 2006 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date August 16, 2007.
- 7. Motor Vehicle Crash Death Rate: number of motor vehicle crash deaths from 2004 to 2006 in age group per 100,000 population in age group.
- 8. Suicide rate: number of suicides from 2004 to 2006 in age group per 100,000 population in age group.
- 9. Rates are based on the National Center for Health Statistics, estimates of the July 1, 2005 United States resident population from the Vintage 2005 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date August 16, 2006.
- 10. Prior to 2005, Idaho BRFSS reported drinking status among all adults (binge drinking = 5 or more drinks in single occasion). Beginning in 2005, Idaho reports gender specific status (5 or more for males, 4 or more for females, in single occasion.